Evolution of "Plants"

Flora & Fauna



Divergence

Seeds & Pollen

Spermatophyta

- Loss of flagella
- Strigolactone DI4 Brassinosteroid

Angiosperms

- Tissue specific specialisation of plastids

Gymnosperms

Tracheophyta

- Vasculature
- Sporophyte dominance
- Della first major duplication event
- Stomata partially lost

Ferns

Lycophyta

Embryophyta Land Plants

- Roots
- Multicellular sporophyte
- Embryogenesis
- Cuticle
- Polyplastidy
- RNA-editing
- Archegonia & antheridia
- 3D-growth
- Phenylpropanoids
- Alternation of multicellular generations
- Mycorrhiza with mycorrhiza
- Whole genome duplication
- Sporopollenin
- YABBY
- KNOX1
- Della
- Stomata

Bryophyta

- Loss
- Reductive evolution

Mosses

- Seta

Liverworts

- YABBY lost
- Stomata lost

Hornworts

- KNOX1 lost

Land Plants

- Penium
- Mesotaenium
- Spirogloea
- WGD

Begin of interaction with substrate microbiome

Zygnematophyceae

- Loss of flagella
- Loss of egg & sperm motility and plasmodesmata

Coleochaetophyceae

- Phragmoplast
- Plasmodesmata
- Apical cell growth
- Hexameric cellulose synthase
- Branching
- Specialised vegetative cells
- Cell-cell interaction
- Biplastidy

Charophyceae

- Egg & sperm
- More elaborate retrograde signaling (GSH) & elaboration of phytohormone signaling
- Cellulose synthase rosettes
- Cytokinin: AHKs, ABA
- Strigolactone transcriptional regulation

Klebsormidiophyceae

- Filamentous growth
- Interaction with microbiota
- Terrestrial stresses

Chlorokybophyceae

- Lateral flagella

Mesostigmatophyceae

- Non-motile vegetative phase
- Adaptation to aerial habitat

Strepto-phyta

- Auxin transport
- Cytokinin
- MIKC-type MADS-box genes
- Dormancy
- Spore priming
- Phytochrome
- Cell wall
- Chlorophyll A/B
- Highly developed chloroplast grana

Chlorophyta

- Loss of land plant-like photorespiration

Prasinodermophyta

- Land plant-like photorespiration

Chloro-plastida

Glauco-phyta

Rhodo-phyta

Proteorhodophytina

Florideophyceae

Bangiophyceae

Cyanidiales

Archae-plastida

Eukaryota

Archaea

Bacteria

Cyanobacteria

- Oxygen Evolving Complex 2400-3700

Endosymbiosis

- Facultative aerobic α-Proteo-bacterium
- Photosynthetic Cyanobacterium
- Plastid
- Mitochondrion

Rhizosphere

- Root Exudates
- Mycorrhiza e.g. Arbuscular Mycorrhizal Fungi

The Terrestrial Habitat

500 Ma

Compared to today:

- Naked rock
- Higher volcanic activity
- O^2 CO^2
- Ozone UV-B

Compared to under water:

- Fluctuating temperatures
- Lack of buoyancy
- Novel pathogens
- Dessicative atmosphere
- High irradiance

How land plants affect climate & oxygenation



Soil formation



Atmospheric evolution



Snowball Earth ~650 Ma

Rodinia ~900 Ma

Drought

Cambrian ~500 Ma

Land water

Anydro-phyta

- Canonical
- aux/iaa TIR auxin signaling
- ARF TF auxin response
- ABA TIR abscisic acid receptor
- GRAS TF plant development

Phragmoplasto-phyta

Late Archean & Early Proterozoic 2.8-2.2 Ga

4.0 Ga Hadean

Phragmoplast



2D & 3D body plan formation



Plasmodesmata



Arabidopsis thaliana

Plastids

Alternation of Generations



Multicellularity among eukaryotes



Global Biomass by taxa